

Monodelphis domestica: a new source of mammalian primary neurons in vitro

Ban, Jelena; Mladinic, Miranda

Source / Izvornik: **Neural Regeneration Research, 2022, 17, 1726 - 1727**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.4103/1673-5374.332139>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:193:519362>

Rights / Prava: [Attribution-NonCommercial-ShareAlike 4.0 International/Imenovanje-Nekomercijalno-Dijeli pod istim uvjetima 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-02-09**

Repository / Repozitorij:



[Repository of the University of Rijeka, Faculty of Biotechnology and Drug Development - BIOTECHRI Repository](#)



Monodelphis domestica: a new source of mammalian primary neurons *in vitro*

Jelena Ban, Miranda Mladinic*

In vitro models have tremendously revolutionized cell biology and biomedical research, reducing the need for *in vivo* experiments, as well as offering the simplified models for easier investigations at molecular and cellular level (for example genetic manipulations, electrophysiological measurements, drug screening etc.). However, the major challenge is to develop long-surviving *in vitro* preparations of post-mitotic cells such as neurons or cardiocytes. To overcome the problem of neuronal inability to proliferate, the immortalized cell lines derived from neuronal tumors have been prepared. Such secondary neuronal cultures are restricted to neuroblastoma-like cells, but their biological relevance is often questionable due to genetic drift and lack of mature, differentiated neuronal phenotypes, which makes primary cultures the better choice. Mammalian central nervous system (CNS) *in vitro* primary cell cultures are mostly prepared from the late embryonic or early postnatal mice and rats. Other mammalian species have been less used, meaning that inter-species diversity is not sufficiently investigated and that the additional comparative analyses are required to avoid misinterpretations in translating the knowledge to humans (Bonfanti and Peretto, 2011).

We have recently established and extensively characterized the long-term primary dissociated cortical neuronal cultures derived from the neonatal grey South American short-tailed opossums [*Monodelphis (M.) domestica*] that we propose as a new source of mammalian CNS cells for *in vitro* developmental and regeneration studies (Petrović et al., 2021). These cultures are unique for the possibility to obtain embryonic-like CNS cells from the postnatal animals, at wide range of postnatal days. For example, the radial glia cells (RGCs), with both neurogenic and gliogenic potential, can be obtained in co-culture with neurons from the cortex of the postnatal day (P) 3–5 opossums and maintained in proliferative conditions for weeks and even months *in vitro*. RGCs cultures can be replated several times, offering a continuous source of new RGCs and *in vitro*-generated neurons. Alternatively, if cultured in serum-free medium, these cells differentiate into almost pure neuronal cultures (more than 90% of neurons in the first few days after plating and more than 80% of neurons during the first two weeks). Furthermore, neuronal cultures can be prepared also from P16–18 opossum cortex that, in addition

to neurons, contain also around 19% of the glial fibrillary acidic protein (GFAP)-positive astrocytes and around 2% of the ionized calcium-binding adaptor molecule 1-positive microglia cells, as well. The percentage of astrocytes is comparable to rat hippocampal neuronal cultures obtained from P2–3 rat hippocampus (around 23%) (Pozzi et al., 2017). The possibility to prepare neuronal cultures from animals at several weeks of postnatal age offers an additional opportunity to investigate specific developmental events and transitions that occur during cortical neurogenesis.

The special characteristics of the opossum CNS cultures come from the fact that opossums are born at a very immature state (Nicholls et al., 1990; Mladinic et al., 2009) which makes them an excellent model for *in vitro* developmental studies. Opossums belong to marsupials, mammalian infraclass diverged from placental mammals around 180 million years ago (Mikkelsen et al., 2007). Their short gestation of 14 days, i.e., 1 week shorter than rodents, allows fast and easy breeding. According to recent developmental transcriptome analysis (Cardoso-Moreira et al., 2019), P3–5 opossums correspond to E15.5–18.5 rat or E14–16 mice embryos, while P16–18 opossums are developmentally similar to neonatal (P1–P2) rat or mice. Their CNS development, that include both neurogenesis and gliogenesis, occurs almost completely postnatally and therefore it offers exceptionally wide time window for developmental studies. In addition, unlike the other marsupials, *M. domestica* lack pouch which makes pups easily accessible. The use of postnatal animals reduces the number of animals needed for studies and reduces their suffering, contributing to the 3Rs (reduce, refine, replace). Shorter gestation and embryonic-like tissue available using postnatal animals with extended postnatal development makes *M. domestica* appealing and ethically improved cell source.

Most importantly, opossums are unique among mammals in their ability to postnatally completely and fully functionally regenerate their spinal cord after injury (Nicholls and Saunders, 1996). This capacity persists until the P12 in the upper cervical part, and until the P17 in the lower, less mature thoracic-lumbar spinal cord segments. The postnatal spinal cord regeneration in opossums has been studied *in vivo* and *in vitro* (Varga et al., 1996) using the intact spinal cord preparation in which the regenerative potential retains the time and space distribution as *in vivo*. However,

both approaches do not allow easy molecular or pharmacological manipulations, unlike cell cultures, which are easier to maintain, manipulate and analyse.

Thus, using developing cortex of P3–5 opossums, we have successfully established long-term and nearly pure neuronal cultures (Petrović et al., 2021). These cultures recapitulate *in vitro* many developmental processes, remarkably similar to what has been observed with well-known rodent models, for example neurite outgrowth, neuronal growth cones (GCs) formation, axon and dendrite specification, synaptogenesis, neurotransmitter synthesis, network formation etc. (Kaech et al., 2012).

For instance, we reported for the first time the efficient *in vitro* generation of neuronal GCs derived from opossum cortex, with an average size of around 40 μm^2 , comparable to what is observed in the rat hippocampal GCs cultured under similar conditions (Pozzi et al., 2017). *M. domestica*-derived GCs are formed at the tips of the growing neurons at the first day *in vitro* (DIV)1 and are identified by actin staining. It will be interesting to further investigate to what extent the cytoskeletal dynamics and expression of receptors for axon guidance molecules, growth factors or neurotrophins that occur in early mammals such as opossums correlate to what is observed in rodents.

Extensive characterization of the cell morphology, composition, differentiation as well as maturation of opossum cortical cultures was done by immunocytochemistry (Petrović et al., 2021). The expression of the neuron-specific β -tubulin III, was confirmed throughout the observation period (up to DIV30) with 4 different β -tubulin III antibodies, having between 90–99.8% sequence similarity between opossum protein and the immunogen of interest. High sequence similarity with rat, mouse or human proteins allows the use of commercially available and widely-used antibodies, such as TUJ1. This marker was used to quantify the percentage of neurons in cultures prepared from two postnatal age groups, P3–5 and P16–18. Both cultures were composed of more than 80% of β -tubulin III-positive neurons during the first week *in vitro*. However, at DIV15 the cultures derived from P16–18 cortex showed significantly lower percentage of neurons (around 55%) and the higher proportion of non-neuronal cells. These observations were comparable to rat hippocampal cultures, in which higher number of glial cells was found in postnatal versus embryonic primary cortical cells. To further study neuronal cells in opossum cortical cultures, two additional neuronal markers were used, namely microtubule-associated protein (MAP) 2 and neuronal nuclei (NeuN, **Figure 1**). Triple staining indicated that the expression of NeuN starts slightly earlier at DIV1, followed by MAP2 and β -tubulin III, with great majority of neurons (more than 86% at DIV1 and more than 93% at DIV4) expressing all




Figure 1 | Neuronal markers expression in nearly pure cortical cultures derived from P5 opossum.

Cells were prepared as described in a previous study (Petrović et al., 2021), fixed at the fourth day *in vitro* and stained for β -tubulin III (TUJ1, red), microtubule-associated protein 2 (MAP2, green), neuronal nuclei (NeuN, magenta) and 4',6-diamidino-2-phenylindole (DAPI) nuclear stain (blue). More than 93% of cells at the fourth day *in vitro* (DIV4) are triple-positive, TUJ1⁺/MAP2⁺/NeuN⁺. Scale bar: 50 μ m. Unpublished data.

three neuronal markers.

To the best of our knowledge, these are the first reports for the expression of MAP2 and NeuN neuronal markers in *M. domestica* primary dissociated cortical cultures. These observations suggest that NeuN can be reliably used as the early neuronal marker and not only as a marker for mature neurons, as it was considered for decades. In addition, we have shown for the first time the expression of synaptic (synapsin) as well as neuronal subtype (VGLUT2 and GAD65) specific markers in *M. domestica*-derived cortical primary cultures, with which is possible to follow neuronal differentiation *in vitro*, as well as the formation of the neuronal networks.

Next, since the gliogenesis starts in opossums around P18, with the morphological switch from bipolar RGCs to multipolar astrocytes (Puzzolo and Mallamaci, 2010), we analysed the expression of glial cell markers (GFAP, vimentin, ionized calcium-binding adaptor molecule 1) in opossum cortical cultures. Our findings are in agreement with the gene expression analysis done by Cardoso et al. (2019), in which developmental correspondences between different mammalian species are reported. We have identified astrocytes and their subtypes, protoplasmic-like and stellate forms, by GFAP and vimentin staining, showing that their morphology was strikingly similar to rodent primary astrocytes, indicating *M. domestica* also as the new source of mammalian astrocytes for *in vitro* investigations. In addition, classification of astrocytes and glial cells in general is still incomplete and it would be interesting to verify if the glial subtypes in opossums resemble more rodent or human-specific glial repertoire.

Perhaps the main advantage that neonatal opossums' cortical cultures offer is the possibility to expand RGCs and generate new neurons and glia *in vitro*. To do that, the cultures can be maintained in specific conditions for several weeks *in vitro*, similar for neuroblastoma cell lines, but with the advantage of having non-transformed, i.e., wild-type neuronal and glial precursor cells in culture, with obvious biological relevance. *M. domestica*-derived stem/progenitor/RGC were identified by their elongated bipolar morphology, GFAP and vimentin expression, proliferative state and

by the potential to form neurospheres. Neurospheres were obtained in suspension, using non-adherent conditions, where they reached the diameters of around 100 μ m and were positive for early progenitor marker paired box gene 2. When replated on adhesive substrate in serum-free conditions, they extended neurites and expressed neuronal markers MAP2 and β -tubulin III. This neurogenic potential offers virtually unlimited *in vitro* source of neurons, particularly useful for stem cell research.

Moreover, the long-term survival of the opossum cortical cultures allowed the formation of the secondary organoid-like structures after several weeks *in vitro*, that consisted of spherical aggregates of neuronal cell bodies from which the thick neurite bundles grew. These results open the possibility to investigate the formation of 3D-cultures combined with 3D biomaterial scaffolds that would more closely mimic the complexity of the CNS *in vivo*.

Thus, the *M. domestica*-derived primary CNS cultures (Petrović et al., 2021) are novel and robust *in vitro* platforms to use in a variety of studies, including development and regeneration. Since the *M. domestica* genome (Mikkelsen et al., 2007; Samollow, 2008) show high degree of protein-coding sequence homology with placental mammals, the opossum cortical cultures represent an excellent new tool to overcome the lacuna in the diversity of the mammalian preparations available for neurobiology studies.

The present work has been conducted on equipment financed by the European Regional Development Fund (ERDF) within the project "Research Infrastructure for Campus-based Laboratories at University of Rijeka" (RC.2.2.06-0001), the Croatian Science Foundation (CSF) grant IP-2016-06-7060, the financial support from the University of Rijeka (18.12.2.1.01, 18-258-6427 and 18-290-1463) and from the International Centre for Genetic Engineering and Biotechnology (ICGEB), Grant/Award Number: CRP/CRO14-03 (to MM).

The authors declare that there is no potential conflict of interest.

Editor note: MM is an Editorial Board member of Neural Regeneration Research. She was blinded from reviewing or making decisions on the manuscript. The article was subject to the journal's standard procedures, with peer review handled independently of this Editorial Board member and their research groups.

Jelena Ban, Miranda Mladinić*

Laboratory for Molecular Neurobiology, Department of Biotechnology, University of Rijeka, Rijeka, Croatia

*Correspondence to: Miranda Mladinić, PhD, mirandamp@biotech.uniri.hr.

https://orcid.org/0000-0002-3985-6629

(Miranda Mladinić)

Date of submission: April 7, 2021

Date of decision: April 19, 2021

Date of acceptance: June 18, 2021

Date of web publication: January 7, 2022

https://doi.org/10.4103/1673-5374.332139

How to cite this article: Ban J, Mladinić M (2022) *Monodelphis domestica*: a new source of mammalian primary neurons *in vitro*. *Neural Regen Res* 17(8):1726-1727.

Open access statement: This is an open access journal, and articles are distributed under the terms of the Creative Commons AttributionNonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

©Article author(s) (unless otherwise stated in the text of the article) 2022. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

References

- Bonfanti L, Peretto P (2011) Adult neurogenesis in mammals—a theme with many variations. *Eur J Neurosci* 34:930-950.
- Cardoso-Moreira M, Halbert J, Vallotton D, Velten B, Chen C, Shao Y, Liechti A, Ascensão K, Rummel C, Ovchinnikova S, Mazin PV, Xenarios I, Harshman K, Mort M, Cooper DN, Sandi C, Soares MJ, Ferreira PG, Afonso S, Carneiro M, et al. (2019) Gene expression across mammalian organ development. *Nature* 571:505-509.
- Kaech S, Huang CF, Banker G (2012) General considerations for live imaging of developing hippocampal neurons in culture. *Cold Spring Harb Protoc* doi:10.1101/pdb.ip068221.
- Mikkelsen TS, Wakefield MJ, Aken B, Amemiya CT, Chang JL, Duke S, Garber M, Gentles AJ, Goodstadt L, Heeger A, Jurka J, Kamal M, Maudsley S, Searle SM, Sharpe T, Baker ML, Batzer MA, Benos PV, Belov K, Clamp M, et al. (2007) Genome of the marsupial *Monodelphis domestica* reveals innovation in non-coding sequences. *Nature* 447:167-177.
- Mladinić M, Müller KJ, Nicholls JG (2009) Central nervous system regeneration: from leech to opossum. *J Physiol* 587:2775-2782.
- Nicholls JG, Stewart RR, Erulkar SD, Saunders NR (1990) Reflexes, fictive respiration and cell division in the brain and spinal cord of the newborn opossum, *Monodelphis domestica*, isolated and maintained *in vitro*. *J Exp Biol* 152:1-15.
- Nicholls J, Saunders N (1996) Regeneration of immature mammalian spinal cord after injury. *Trends Neurosci* 19:229-234.
- Petrović A, Ban J, Tomljanović I, Pongrac M, Ivaničić M, Mikšašević S, Mladinić M (2021) Establishment of long-term primary cortical neuronal cultures from neonatal opossum *Monodelphis domestica*. *Front Cell Neurosci* doi:10.3389/fncel.2021.661492.
- Pozzi D, Ban J, Iseppon F, Torre V (2017) An improved method for growing neurons: comparison with standard protocols. *J Neurosci Methods* 280:1-10.
- Puzzolo E, Mallamaci A (2010) Cortico-cerebral histogenesis in the opossum *Monodelphis domestica*: generation of a hexalaminar neocortex in the absence of a basal proliferative compartment. *Neural Dev* 5:8.
- Samollow PB (2008) The opossum genome: insights and opportunities from an alternative mammal. *Genome Res* 18:1199-1215.
- Varga ZM, Fernandez J, Blackshaw S, Martin AR, Müller KJ, Adams WB, Nicholls JG (1996) Neurite outgrowth through lesions of neonatal opossum spinal cord *in culture*. *J Comp Neurol* 366:600-612.

C-Editors: Zhao M, Liu WJ, Qiu Y; T-Editor: Jia Y